



Volume 162 · Number 3 · March 2010

PHYSIOLOGICAL ECOLOGY

Activation of the immune system promotes insect dispersal in the wild  
J. Suhonen · J. Horkkari · M. J. Rantala 541

Why eat? Trait and habitat correlates of variation in germination speed among *Kalanchoe* annuals  
M. Kos · P. Paschold 549

Sex-specific plant responses to light intensity and canopy openness: implications for spatial segregation of the sexes  
K.E. Green · C.R. Sogho · P.H. Crowley · D.N. McLetchie 561

The impact of protein quality on stable nitrogen isotope ratio discrimination and assimilated diet estimation  
C.T. Robbins · L.A. Falcoff · S.T. Florin 571

Deviation from strict homeostasis across multiple trophic levels in an invertebrate consumer assemblage exposed to high chronic phosphorus enrichment in a Neotropical stream  
G.E. Small · C.M. Pingle 581

BEHAVIORAL ECOLOGY

Family movements before independence influence natal dispersal in a territorial songbird  
E. Matthysen · T. Van Overveld · T. Van de Casteele · F. Adriansen 591

What are the evolutionary constraints on larval growth in a trophically transmitted parasite?  
D.P. Bensch 599

Is spatial autocorrelation an intrinsic property of territory size?  
M. Valcarlos · B. Kempenaers 609

Males make poor meals: a comparison of nutrient extraction during sexual cannibalism and predation  
S.M. Wilker · A.L. Rypstra 617

Observational evidence of risk-sensitive reproductive allocation in a long-lived mammal  
B.-J. Bilde · T. Tveraa · P. Fausch · K. Langeland 627

POPULATION ECOLOGY

Population synchrony of a native fish across three Laurentian Great Lakes: evaluating the effects of dispersal and climate  
D.B. Bunnell · J.V. Adams · G.T. Gorman · C.P. Madenjian · S.C. Riley · E.F. Roseman · J.S. Schaeffer 641

Individual growth rates in natural field vole, *Microtus agrestis*, populations exhibiting cyclic population dynamics  
S.J. Barber · K. Lindén · S. Teller · A. Douglas · P. Bekasman · A. Smith · M. Begon 653

Offspring size and timing of hatching determine survival and reproductive output in a lizard  
T. Liljer · M. Olsson 663

PLANT-ANIMAL INTERACTIONS

Developmental plasticity and reduced susceptibility to natural enemies following host plant defoliation in a specialized herbivore  
G.R. Hood · J.R. De 673

Induced resistance in a brown edge: phlorotannins, genotypic variation and fitness costs for the crustacean herbivore  
F. Haavisto · T. Vähkangas · V. Jormalainen 685

Tree squirrel habitat selection and predispersal seed predation in a declining subalpine conifer  
S.T. McKinney · C.E. Feder 697

COMMUNITY ECOLOGY

Your worst enemy could be your best friend: predator contributions to invasion resistance and persistence of natives  
S.A. Juliano · L.P. Lounibos · N. Nishimura · K. Greene 709

Woody-grass ratios in a grassy arid system are limited by multi-causal interactions of abiotic constraint, competition and fire  
C.E.M. Naro · P.J. Clarke 719

Interaction intensity and importance along two stress gradients: adding shape to the stress-gradient hypothesis  
P.C. Le Roux · M.A. McGeoch 733

A non-trophic interaction chain links predators in different spatial niches  
B.P. Pissard · W.E. Snyder 747

ECOSYSTEM ECOLOGY

Natural enemy diversity reduces temporal variability in wasp but not bee parasitism  
D. Veddeler · J. Tylianakis · T. Tscharntke · A.-M. Klein 755

Biodiversity at the plant-soil interface: microbial abundance and community structure respond to litter mixing  
S.K. Chapman · G.S. Newman 763

Litter decomposition and nutrient release as affected by soil nitrogen availability and litter quality in a semi-arid grassland ecosystem  
P. Liu · J. Huang · D.J. Sun · X. Han 771

Leaf litter traits of invasive species slow down decomposition compared to Spanish natives: a broad phylogenetic comparison  
D. Gaddy · P. Castro-Díez · R.S.P. Van Logtestijn · J.H.C. Cornelissen · F. Valladares 781

GLOBAL CHANGE ECOLOGY

Elevated carbon dioxide alters impacts of precipitation pulses on ecosystem photosynthesis and respiration in a semi-arid grassland  
S. Baatman · J.L. Hester-White · E. Pienaar · D.G. Williams · J.A. Morgan · J. Newcomb 791

CONSERVATION ECOLOGY

Linking snake habitat use to nest predation risk in grassland birds: the dangers of shrub cover  
P.E. Rlug · S.L. Jackel · K.A. With 803



This article was published in the above mentioned Springer issue.  
The material, including all portions thereof, is protected by copyright;  
all rights are held exclusively by Springer Science + Business Media.  
The material is for personal use only;  
commercial use is not permitted.  
Unauthorized reproduction, transfer and/or use  
may be a violation of criminal as well as civil law.

# Leaf litter traits of invasive species slow down decomposition compared to Spanish natives: a broad phylogenetic comparison

Oscar Godoy · Pilar Castro-Díez ·  
Richard S. P. Van Logtestijn ·  
Johannes H. C. Cornelissen · Fernando Valladares

Received: 28 January 2009 / Accepted: 2 November 2009 / Published online: 22 November 2009  
© Springer-Verlag 2009

**Abstract** Leaf traits related to the performance of invasive alien species can influence nutrient cycling through litter decomposition. However, there is no consensus yet about whether there are consistent differences in functional leaf traits between invasive and native species that also manifest themselves through their “after life” effects on litter decomposition. When addressing this question it is

important to avoid confounding effects of other plant traits related to early phylogenetic divergences and to understand the mechanism underlying the observed results to predict which invasive species will exert larger effects on nutrient cycling. We compared initial leaf litter traits, and their effect on decomposability as tested in standardized incubations, in 19 invasive-native pairs of co-familial species from Spain. They included 12 woody and seven herbaceous alien species representative of the Spanish invasive flora. The predictive power of leaf litter decomposition rates followed the order: growth form > family > status (invasive vs. native) > leaf type. Within species pairs litter decomposition tended to be slower and more dependent on N and P in invaders than in natives. This difference was likely driven by the higher lignin content of invader leaves. Although our study has the limitation of not representing the natural conditions from each invaded community, it suggests a potential slowing down of the nutrient cycle at ecosystem scale upon invasion.

Communicated by Stephan Hattenschwiler.

**Electronic supplementary material** The online version of this article (doi:10.1007/s00442-009-1512-9) contains supplementary material, which is available to authorized users.

O. Godoy (✉) · F. Valladares  
Instituto de Recursos Naturales,  
Centro de Ciencias Medioambientales, CSIC,  
Serrano 115 dpdo, 28006 Madrid, Spain  
e-mail: ogodoy@ccma.csic.es

O. Godoy · P. Castro-Díez  
Departamento de Ecología, Facultad de Ciencias,  
Universidad de Alcalá, 28871 Alcalá de Henares, Madrid, Spain

R. S. P. Van Logtestijn · J. H. C. Cornelissen  
Department of Systems Ecology,  
Faculty of Earth and Life Sciences, Institute of Ecological Science,  
Vrije Universiteit, De Boelelaan 1085, 1081 HV Amsterdam,  
The Netherlands

F. Valladares  
Departamento de Biología y Geología,  
Área de Biodiversidad & Conservación,  
Universidad Rey Juan Carlos, ESCET,  
Tulipán s/n, 28933 Móstoles, Madrid, Spain

F. Valladares  
Laboratorio Internacional de Cambio Global (LINCglobal),  
Departamento de Ecología, Facultad de Ciencias Biológicas,  
PUC, UC-CSIC, Alameda 340, 6513677 Santiago, Chile

**Keywords** Nitrogen · Phosphorus · Lignin · Calcium · Phylogenetically independent contrast

## Introduction

Litter decomposition is a key process in nutrient cycling (Swift et al. 1979; Hobbie 1992; Sayer 2006). Rates of litter decomposition can greatly influence nutrient availability and thereby other ecosystem processes (Vitousek and Howarth 1991). At the community level, decomposition rates are controlled by species or functional type composition (Hättenschwiler and Vitousek 2000; Garnier et al. 2004; Quested et al. 2005). Therefore, the introduction or invasion of alien species into new ecosystems can cause

profound changes in community structure and function (Mack et al. 2000; Kolar and Lodge 2001), including changes in litter decomposition and nutrient cycling (Allison and Vitousek 2004; Ashton et al. 2005). To determine the direction of these changes, several studies have compared litter decomposition between invasive alien species and natives. However, results differ greatly between studies, taxa, and regions (Ehrenfeld 2003) and no generalization (a frequent objective in the study of biological invasion) has arisen so far. Faster decomposition of litter from invasive alien species compared to native litter has been reported when invaders were N<sub>2</sub>-fixers and native species were not (Liao et al. 2008), when nutrient concentration in the exotic litter was higher than in the native one (Allison and Vitousek 2004) and when the specific leaf area (SLA) of invaders was higher than that of native species (Castro-Díez et al. 2009; Kueffer et al. 2008). By contrast, slower decomposition of invasive alien species' leaf litter was found when it had a higher polyphenolic content (Knight et al. 2007), higher lignin content or higher C/N ratio than native litter (Drenovsky and Batten 2007).

Differences between studies may be partly due to overrepresentation of particular regions (e.g. Hawaii islands; Allison and Vitousek 2004) or functional types (e.g. at least 1/3 of the articles involve N<sub>2</sub>-fixers; Levine et al. 2003), and they have generally included only a few species pairs, which have not accounted for phylogenetic and life history differences. Comparing large numbers of invasive and native species by means of phylogenetic independent contrasts is necessary to unveil general trends whilst avoiding confounding effects of phylogeny.

As important as avoiding confounding effects, is to understand the mechanisms underlying the direction of the change upon invasion. This will potentially help to predict which invasive alien species are likely to exert larger effects on nutrient cycling and which communities are likely to suffer larger impacts (Levine et al. 2003). Invasive species possessing leaf litter traits similar to natives will be functionally redundant in the nutrient cycling mechanisms and no changes may be expected. By contrast, invasive species with leaf litter traits largely different from those of natives, may impact the nutrient cycling. Thus, the direction and magnitude of the change will depend on the trait difference between these groups (Dukes and Mooney 2004; Strauss et al. 2006). Under Mediterranean soil conditions, where the availability of N and P is low (Gallardo and Parama 2007), we may predict that leaf litter from invasive alien species with traits that retard decomposition will inhibit and slow down nutrient cycling, while those promoting decomposition will have no effect (because of microorganism immobilization in cases of low nutrient soil concentration) or will slightly speed up nutrient cycling (Knorr et al. 2005).

By using this less frequently employed approach, we have the additional opportunity to link ecosystem impacts to functional strategies of invasive species, since many leaf litter traits are directly related to plant performance strategies such as competitive ability (e.g. variation in N content), defence against herbivores (e.g. variation in polyphenolic compounds) or resource use efficiency (e.g. variation in lignin content). Thus, while many studies have tried to link species traits with invasiveness, the importance of studying these traits is more related to differences in ecosystem impacts than in invasiveness.

The objective of the present study was to compare leaf litter decomposability between invasive alien and native species from Mediterranean ecosystems in Spain by means of an experimental design using phylogenetically independent contrasts (Armstrong and Westoby 1993). We addressed the following questions:

1. Do leaf litter properties of invasive alien species differ consistently from those of native ones?
2. Does leaf litter of invasive alien species decompose faster than that of native species?
3. Do initial leaf litter traits influence the decomposition process in the same way in invasive and native species?

To address these questions, a set of 19 invasive-native co-familial pairs were selected and subjected to an experimental standardized screening for leaf litter decomposability together with physical–chemical analyses.

## Materials and methods

### Species selection

Our species set comprised introduced species that are clearly invasive (Pyšek et al. 2004), local dominants in some native ecosystems (Valéry et al. 2008) and with potential impact in the invaded ecosystems (Richardson et al. 2000). Species compiled by *The Atlas of the Invasive Alien Species in Spain* (Sanz Elorza et al. 2004) matched our criteria of invasive species definition. Thus, it was used to select the invasive alien species and the localities where the senesced leaf material was collected, ranging from central to eastern Spain (Table 1). The final selection of invasive species was done on the basis of leaf litter availability and aimed to represent of a broad diversity of growth forms (annual and biannual herbs, lianas, shrubs, and trees), invaded habitats (woodlands, shrublands, grasslands, and riparian areas), and different life histories. Nomenclature followed the Iberian Flora (Castroviejo 1986–2008). Each of the selected species was paired with one closely related native species based on two criteria: (1) within each pair, the native species had to belong to the same family as the

**Table 1** Invasive-native phylogenetic pairs, type of ecological community invaded and locations selected. *W* Woody species, *H* herbaceous species, *E* evergreen species, *D* deciduous species

Family	Invasive species	Native species	Ecological community	W/H	E/D	Spatial location, Spanish province
Aceraceae	<i>Acer negundo</i>	<i>Acer campestre</i>	Woodland	W	D	40°30'N, 3°20'W, Madrid
Anacardiaceae	<i>Schinus molle</i>	<i>Pistacia terebinthus</i>	Shrubland	W	E/D	39°10'N, 0°16'W, Valencia/ 40°34'N, 3°09'W, Guadalajara
Asteraceae	<i>Achillea filipendula</i>	<i>Achillea millefolium</i>	Grassland	H	D	40°30'N, 3°20'W, Madrid
Caprifoliaceae	<i>Lonicera japonica</i>	<i>Lonicera etrusca</i>	Shrubland and riparian areas	H	D	40°30'N, 3°20'W, Madrid
Amaranthaceae	<i>Atriplex semibaccata</i>	<i>Atriplex halimus</i>	Shrubland	H	E	38°14'N 0°32'W, Alicante/ 40°30'N, 3°20'W, Madrid
Convolvulaceae	<i>Ipomoea indica</i>	<i>Calystegia sepium</i>	Riparian areas	H	D	38°24'N 0°26'W, Alicante
Elaeagnaceae/ Rhamnaceae	<i>Elaeagnus angustifolia</i>	<i>Rhamnus alaternus</i>	Shrubland	W	D/E	40°30'N, 3°20'W, Madrid
Fabaceae	<i>Gleditsia triacanthos</i>	<i>Colutea arborescens</i>	Shrubland	W	D	40°34'N, 3°09'W, Guadalajara
Fabaceae	<i>Sophora japonica</i>	<i>Ceratonia siliqua</i>	Woodland	W	E	40°30'N, 3°20'W, Madrid/ 38°33'N 0°31'W, Alicante
Fabaceae	<i>Robinia pseudoacacia</i>	<i>Coronilla glauca</i>	Shrubland	W	D/E	40°30'N, 3°20'W, Madrid
Fabaceae	<i>Acacia saligna</i>	<i>Anagyris foetida</i>	Shrubland	W	E/D	38°31'N 0°10'W, Alicante/ 39°09'N 0°21'W, Valencia
Myrtaceae	<i>Eucalyptus globulus</i>	<i>Myrtus communis</i>	Shrubland	W	E	40°30'N, 3°20'W, Madrid
Onagraceae	<i>Oenothera biennis</i>	<i>Epilobium hirsutum</i>	Grasslands and riparian areas	H	D	39°06'N 0°23'W, Valencia/ 40°30'N, 3°20'W, Madrid
Pinaceae	<i>Pinus radiata</i>	<i>Pinus pinaster</i>	Woodland	W	E	40°30'N, 3°20'W, Madrid
Simaroubaceae/ Cneoraceae	<i>Ailanthus altissima</i>	<i>Cneorum tricoccom</i>	Shrubland	W	D/E	40°30'N, 3°20'W, Madrid
Solanaceae	<i>Nicotiana glauca</i>	<i>Lycium intricatum</i>	Shrubland	H	E	38°05'N 0°40'W, Alicante
Solanaceae	<i>Solanum bonaeriense</i>	<i>Solanum nigrum</i>	Shrubland and grassland	H	D	39°33'N 2°42'W, Mallorca/ 40°30'N, 3°20'W, Madrid
Ulmaceae	<i>Ulmus pumila</i>	<i>Ulmus minor</i>	Shrubland and riparian areas	W	D	40°28'N, 3°23'W, Madrid
Verbenaceae	<i>Lantana camara</i>	<i>Vitex agnus-castus</i>	Shrubland	W	E/D	38°30'N, 0°14'W, Alicante

invasive species; and (2) the two species had to belong to the same growth form (i.e. herbaceous vs. woody species).

The selection of phylogenetically independent invasive-native pairs for methodological reasons may induce biases in the ecological meaning of our results. To minimize this problem, we conducted a screening to select native species with a common occurrence throughout Spain and especially in the eastern part, which hosts the highest diversity and abundance of invasive alien species (Gasso et al. 2009). Native species with a small distribution range, and those which are rare or endangered were excluded. Additionally, we discarded those native species occurring in habitats different from those where alien species invade or with occurrence in successional community stages different from natives. The extended herbarium database at the Complutense University of Madrid was used to select, within pairs, native species recorded in the same localities or next to where the alien species pair invade. This procedure has permitted us to retain the potential ecological meaning of our

results in the whole range of the invasive species, except for the case of *Ailanthus altissima* vs. *Cneorum tricoccom* (Table 1), which only has ecological sense in the Balearic Islands and north of Cataluña.

#### Collection of leaf litter

Recently senesced, undecomposed and herbivore-damage-free leaf litters of invasive alien species and their native relatives were collected following the criteria of Cornelissen (1996). Leaf litter from deciduous woody species was collected throughout the autumn 2005, while leaf litter of evergreen woody and herbaceous species was collected in the spring 2006, according to their natural leaf senescence period. Ten mature individuals for each species of the same population at full-sun exposure (in the case of trees) were preferentially chosen for senesced leaf collection. Large variation in the proportion of petioles to lamina was found between species ( $6.7 \pm 5.3\%$ ), but petioles from all species

were discarded to assess the decomposition ability of leaf litter, since the variability within pairs was low ( $1.2 \pm 0.4\%$ ). Approximately 40 g of senesced leaves per species was obtained after pooling the leaves from all individuals. They were transported to the laboratory, air-dried at 20–25°C and stored in open paper bags until further analysis.

#### Litter decomposition experiment set up

For each species, four subsamples of  $1.0 \pm 0.1$  g air-dried senesced leaves were used to calculate their initial water content by weighing them again after 3 days in the oven at 60°C. Sixteen samples of air-dried leaves were extracted from each species pool, weighed ( $1.0 \pm 0.1$  g) and corrected for water content. Then each of these samples was sealed into a 1-mm-mesh nylon bag to allow free movement of macro- and meso-decomposers into the bags (Cornelissen 1996). Large leaf size variation between species lead to variation in bag size in order to spread out whole leaves in one single layer, since surface access to soil decomposers is a determinant of decomposition rate (Cornelissen 1996). Litter bags were buried randomly in a purpose-built litter decomposition bed (7 m  $\times$  3.5 m) at Alcalá University Botanical Garden (Madrid province, Central Spain). This bed consisted of a thoroughly mixed 5-cm litter layer collected from a nearby mature *Quercus* spp. forest. We selected this material trying to create an incubation matrix of natural non-invaded forest communities representative of Spain with close-to-natural microbial and soil fauna activity. Additionally, we covered the bed with a single layer of senesced *Platanus hispanica* leaves. The decomposition bed was displayed below a shade frame covered by a neutral shade mesh, which removed 30% of the solar radiation, in order to avoid rapid water loss, especially in summer. On 23 December 2005, the litter bags of the autumn-deciduous invasive-native pairs (pair numbers 1, 4, 8 and 18; Table 1) were placed in the litter decomposition bed at 2-cm depth in the matrix, while the rest of the 15 pairs (where at least one of the species pair shed its leaves in spring) were placed there on 22 June 2006. Two harvests of eight litterbags per species were collected, the first after 147 and 143 days of incubation (for the December and June pairs, respectively), and the final after 289 and 231 days. The first harvest date was chosen by expert criteria to select a steady-state of mean litter decomposition around 50% of mass loss taking into consideration all species, while the final harvest date was chosen when the species with faster litter decomposition (*Colutea arborescens*, *Anagyris foetida*) were close to full decomposition. Eight additional litterbags of *Acer campestre*, *Lonicera japonica* and *Ulmus pumila* were placed in each of the two periods of our experiment to assess the effect of starting date on the decomposition

rate. At each harvest, the material remaining in each litter-bag was carefully brushed clean, dried at 60°C during 3 days, and weighed to calculate the percentage of litter mass loss. For contamination by dust particles see below.

At the beginning of the experiment, three 2.5  $\times$  2.5-cm anion and two 2.5  $\times$  2.5-cm cation exchange resin membranes (types I-100 and I-200; Electropure excellion, Laguna Hills, Calif.) fixed to a plastic label were introduced into the soil at three equidistant points of the litter bed at 4-cm depth, and left for 15 days to evaluate inorganic N and P availability (Abrams and Jarrell 1992; Subler et al. 1995). Resin membranes had been previously conditioned in the lab by immersing them in demineralised water at 82–90°C for 48 h. Ions were extracted from the resins using 20–25 ml KCl and ammonia, nitrate and phosphate contents were assessed following Sims et al. (1995). Finally, local air temperature, available photon flux density (PPFD) and soil moisture within the first 15-cm soil depth were registered every 15 min during the litter decomposition experiment with a data logger (HOBO model H08-006-04; Onset, Pocasset, Mass.) and measured by external sensors [LI-COR, Lincoln, Neb.; ThetaProbe sensors (Delta-T Devices, Cambridge, UK)]. The litter decomposition bed was occasionally watered during dry periods, when soil moisture dropped below 60% gravimetric water content (GWC).

#### Measurements

The initial area of the senesced leaves was measured in eight subsamples of eight leaves each per species (leaf area meter device; Delta-T), which were then oven-dried at 60°C for 3 days and weighed to calculate their SLA (= leaf area/leaf dry mass). The remaining leaf litter, after removing samples for litter decomposition and SLA assessment, was ground in a single pool per species, using a Cullati mill (1-mm particle size). Chemical analyses to determine litter leaf traits were conducted at the laboratory of the Department of Systems Ecology, Vrije Universiteit, Amsterdam, The Netherlands. Total initial concentrations of organic C, N, P, Ca, K, Mg and pH were calculated from a set of four ground subsamples per species, while initial lignin and cellulose contents were assessed in one ground sample per species. C and N were also measured in four ground samples of the final harvest. Finally, in four subsamples per species, ash content was measured (mass loss on ignition, 575°C) to correct litter sample dry mass for possible sand or clay contamination. Since some plants can contain significant amounts of silica and other minerals in their tissues (Epstein 1994), we only corrected for ash fractions greater than 2–3% for species where we observed or suspected contamination by dust.

Total C and N were determined by dry combustion with a NA1500 series II elemental analyser (Carlo Erba,

Rodana, Italy). After digestion (of ground leaf material) in a 1:4 mixture of 37% (v/v) HCl and 65% (v/v) HNO<sub>3</sub> (Sneller et al. 1999) P was measured colorimetrically, Ca and Mg were measured with AAS and K with AES (both: Perkin Elmer 1100, Waltham, Mass.). pH was measured using a narrow (5 mm diameter) SenTix Mic electrode connected to an Inolab Level 2 pH meter (both: WTW, Weilheim, Germany). Initial litter lignin and cellulose concentration was determined following Poorter and Villar (1997): in short, after several extraction steps to ensure that only cellulose and lignin made up the composition of the residue of the sample, the C and N concentrations of this residue were used to calculate the lignin concentration, based upon the difference in C content between cellulose and lignin.

### Statistics

We applied Statistica 6.1 (StatSoft) and R package (library “stats”; R Development Core Team 2009) for all analyses below. Percentage of litter mass loss at the first and final harvests (with initial mass as the reference for both harvests) was arcsine square-root( $\times/100$ ) transformed to approach normal distributions and homogeneity of variance (Quested et al. 2003). Given that some families were over-represented in our dataset (e.g. Fabaceae; Table 1) and that the 19 species pairs were not randomly selected, a Mantel test was conducted to test whether differences in litter decomposition within pairs were influenced by pair selection. The phylogenetic distance from one species to another between all the species pairs was calculated through the first common ancestor to both species using the plant phylogenetic supertree described by Soltis et al. (2000) and modifications by Bremer et al. (2003).

Pairwise Student *t*-tests compared initial physical–chemical leaf litter traits and percentage of litter mass loss across all pairs of invasive alien species and the native co-familial counterpart, at the first and final harvests, respectively. The effect of the start date of incubation (autumn or spring) on the percentage of litter mass loss was assessed for the three control species (*A. campestre*, *L. japonica* and *U. pumila*) by a two-way ANOVA using time and species as main independent factors.

In order to assess the relative importance of different predictors for the litter mass loss, a four-way ANOVA was conducted taking as the main independent variables, Growth form, Family nested in Growth form—because none of the families comprised both woody and herbaceous species—Leaf type (deciduous or evergreen) and Plant status (invasive or native). Additionally, multiple-regression models were conducted to evaluate which senesced leaf traits better predicted the percentage of litter mass loss in each species group (invasive alien and native species) at each harvest. Pearson correlations were conducted to check

for co-linearity between predictors. Predictors which were not correlated or were poorly correlated were chosen ( $r < 0.10$ ), while among those highly correlated ( $r > 0.70$ ) only one predictor with a high ecological meaning was selected (Fowler and Cohen 1992). Therefore, initial lignin concentration was discarded because it was highly correlated with the initial C concentration in both species sets (invasive species  $r > 0.80$  and native species  $r > 0.75$ ). Different models were fitted selecting different sets of predictive variables by the approach based on information theory (Akaike information criteria; AIC; Burnham and Anderson 2002). AIC was calculated for each model and used to choose the best fitted. A value of difference in AIC ( $\Delta$ AIC) higher than 10 represents a poorly fitted model compared to the best model, whereas a value of less than 2 indicates that both models are equivalent (Burnham and Anderson 2002).

### Results

#### Litter bed characteristics

The most abundant form of available mineral N in the litter decomposition bed was NO<sub>3</sub><sup>-</sup>-N (1.82 μg m<sup>-2</sup> day<sup>-1</sup>), which was two times more than NH<sub>4</sub><sup>+</sup>-N (0.86 μg m<sup>-2</sup> day<sup>-1</sup>). The availability of PO<sub>4</sub><sup>-</sup>-P was 0.72 μg m<sup>-2</sup> day<sup>-1</sup>, and thus the N/P ratio of the available pool was 1.9. Mean ( $\pm$ SE) air temperature, soil GWC and daily PPFD during the first incubation period for deciduous autumn pairs (see “Materials and methods”) were 10.21 ( $\pm$ 7.06)°C, 68.74 ( $\pm$ 3.46)% and 24 mol m<sup>-2</sup> day<sup>-1</sup>, respectively, whereas for the final incubation period with the rest of the pairs the, respective, means were 16.84 ( $\pm$ 9.77)°C, 65.12 ( $\pm$ 5.73)% and 32 mol m<sup>-2</sup> day<sup>-1</sup>.

#### Leaf litter traits, decomposition rates and decomposition predictors

SLA did not differ between invasive alien and native species (Table 2). Initial Ca and lignin concentrations were significantly higher in invasive species. Total C concentration was marginally higher in invasive species. Cumulative mass loss at the final harvest was significantly higher in invasive species of a pair than in their co-familial native species, whereas a similar trend was not significant at the first harvest (Table 2). Mean values of all measured traits for each species are shown in the Electronic supplementary material (Table S1). Mantel test R (% mass loss – phylogeny = 1.34;  $P = 0.529$ ), showed that differences in % litter mass loss between invasive and native species were not influenced by the phylogenetic structure of the invasive-native pair selection. In addition,

**Table 2** Leaf litter trait comparison using a pairwise *t*-test based on the 19 invasive-native co-familial pairs. Mean values  $\pm$  SE are shown. Each pair was considered a replicate

Leaf litter traits	Invasive	Native	<i>t</i> -value	<i>P</i>
% Litter mass first harvest	0.36 $\pm$ 0.01	0.42 $\pm$ 0.01	-1.84	0.11
% Litter mass final harvest	0.52 $\pm$ 0.02	0.60 $\pm$ 0.02	-2.02	0.049*
Initial N (%)	1.36 $\pm$ 0.01	1.38 $\pm$ 0.01	-0.13	0.90
Initial C (%)	51.87 $\pm$ 1.98	50.98 $\pm$ 2.10	1.79	0.091
Initial C:N	50.25 $\pm$ 2.67	47.19 $\pm$ 2.36	0.47	0.65
Final N (%)	1.95 $\pm$ 0.39	2.04 $\pm$ 0.63	-0.52	0.62
Final C (%)	52.68 $\pm$ 1.68	52.00 $\pm$ 1.15	0.95	0.21
Final C:N	35.36 $\pm$ 3.04	32.77 $\pm$ 3.12	0.63	0.55
Initial P (mg/g)	0.13 $\pm$ 0.08	0.16 $\pm$ 0.07	-1.03	0.31
Initial Ca (mg/g)	4.10 $\pm$ 0.03	3.21 $\pm$ 0.02	2.17	0.041*
Initial K (mg/g)	1.37 $\pm$ 0.01	1.34 $\pm$ 0.01	0.08	0.95
Initial Mg (mg/g)	0.72 $\pm$ 0.05	0.72 $\pm$ 0.04	-0.01	0.99
Total initial cations (mg/g)	5.86 $\pm$ 0.05	5.27 $\pm$ 0.05	1.48	0.16
pH	5.89 $\pm$ 1.13	5.55 $\pm$ 0.61	1.55	0.25
Initial lignin (mg/g)	139.39 $\pm$ 2.92	95.19 $\pm$ 1.53	2.61	0.029*
Initial cellulose (mg/g)	166.10 $\pm$ 1.92	170.39 $\pm$ 2.04	-0.32	0.68
SLA (m <sup>2</sup> /kg)	9.87 $\pm$ 3.54	9.32 $\pm$ 2.92	-0.65	0.52


\* *P* < 0.05 (between pairs)

two-way ANOVA showed that the effects of the start date of incubation were comparable because significant differences were only found for species ( $F_{2,90} = 41.68$ ,  $P < 0.001$ ), but not for time period ( $F_{1,90} = 0.09$ ,  $P = 0.769$ ) or the time period by species interaction ( $F_{2,90} = 1.66$ ,  $P = 0.206$ ).

Four-way ANOVA showed that growth form was the most important categorical factor explaining the variation of final litter mass loss across the whole data set, mass loss being faster in herbaceous than in woody species. Family was the second best predictor, followed by plant status (invasive-native), native species decompose faster than invasive species, and finally leaf type, deciduous decomposing faster than evergreen species. No significant effect was found in the interaction between plant status and the other categorical factors (Fig. 1).

#### Leaf litter decomposition processes

Total initial concentrations of C, N, P, cations (sum of Ca + K + Mg), pH, cellulose and SLA were the independent predictors selected to assess the relative importance of leaf litter traits in litter decomposition of both harvests and for the invasive and native species groups separately (Table 3). Initial lignin was excluded due to its high correlation with initial C concentration (see “Materials and methods”). Multiple-regression models revealed that the predictors were different between invasive and native species and between harvests (Table 3). N was related positively to litter decomposition in all cases and its relative importance increased at the final harvest for invasive



**Fig. 1** Four-way ANOVA assigning the relative importance of Family, Growth form (woody/herbaceous), Leaf type (evergreen/deciduous) and Status (invasive/native) on percentage mass of leaf litter, based on the 19 invasive-native co-familial pairs (see Table 1). Mass loss % was arcsine [square-root( $\times/100$ )] transformed prior to analysis. Family, as a predictor, was nested in Growth form because there was no Family with both woody and herbaceous species. Explained/residual variance corresponds with the *F*-values of the ANOVA

species. P was the most important positive predictor of % litter mass loss, but only for invasive species at the final harvest. C and thus lignin were negatively related to litter mass loss in native species across the whole incubation period, while only in the first harvest in invasive species (Table 3). pH was a positive predictor of native litter decomposition rate at the final harvest and of invasive species only at the first harvest. A positive relation between decomposition rate and SLA only emerged at final harvest for both groups of species. Finally, cellulose was slightly

**Table 3** Multiple-regression models<sup>a</sup> for invasive and native species at first and final harvest. Dependent variable was the decomposition rate within harvest and independent predictors<sup>b</sup> were selected among initial leaf traits. SLA Specific leaf area

Group	C <sup>c</sup>	N	P	Cations	pH	SLA	Cellulose	AIC	Δ AIC	r <sup>2</sup>
Invasive first harvest	-0.08*	0.07*						-62.028	0.0	0.21
	-0.08*	0.06*			0.12*			-61.434	0.6	0.22
	-0.09*				0.13*			-60.652	1.4	0.18
	-0.08*	0.07*					-0.01	-60.405	1.6	0.19
	-0.08*	0.06*		0.01				-60.211	1.8	0.20
	-0.09*	0.07*				0.04		-60.208	1.8	0.17
Native first harvest	-0.23*	0.12*		0.02*	0.04	0.05	-0.06*	-118.058	0.0	0.73
	-0.22*	0.11*		0.02*	0.05		-0.06*	-117.004	1.1	0.74
Invasive final harvest	-0.06	0.33*	0.65*	0.03*		0.09		-56.183	0	0.52
	-0.06	0.33*	0.66*	0.04*	-0.02	0.11*		-56.031	0.2	0.52
	-0.06	0.32*	0.66*	0.03*		0.01	0.03	-55.165	1	0.52
	-0.06	0.33*	0.67*	0.04*	-0.02	0.11*	0.03	-54.655	1.5	0.51
	-0.07	0.35*		0.04*	-0.02	0.11*		-54.574	1.6	0.47
Native final harvest	-0.23*	0.11*		0.02*	0.18*	0.14*	-0.10*	-75.564	0.0	0.69
	-0.23*	0.12*	-0.12	0.02*	0.19*	0.15*	-0.11*	-73.856	1.7	0.70

\*  $P < 0.05$  (significant in the fitted model)

<sup>a</sup> Models are ranked in each case by Akaike's information criteria (AIC) from best to worst-fitting model, and only the models with  $\Delta AIC < 2$  are presented. For each variable entering the model the standardized regression coefficient is shown to evaluate its relative importance. Empty cells indicate that the leaf trait was not included in the model

<sup>b</sup> Predictor variables: C (%), N (%), P (%), cation [Mg (%) + Ca (%) + K (%)] in litter before incubation

<sup>c</sup> Including initial lignin concentration

negatively related to decomposition rate in native species at both harvests (Table 3).

The goodness-of-fit of the models ( $r^2$ ) was higher for native than for invasive species. Regression models generally predicted litter decomposition rate better at the final than at the first harvest for invasive species (Table 3).

## Discussion

### Leaf litter characteristics and mass loss rates

Leaf litter of alien species invading Mediterranean ecosystems in Spain tended to decompose slightly more slowly than that of the co-familial native species. This could be due to a large initial lignin content, which was significantly higher in invasive alien species across invasive-native species pairs. This well-studied relationship (Swift et al. 1979) was recently confirmed within multiple sites worldwide (Cornwell et al. 2008). Invasive alien species combine higher lignin with higher Ca initial content (Table 2). Ca plays an important role in lignin synthesis and a deficit of this macroelement in soils limits lignification of plant tissues (Lautner et al. 2007). Indeed, the soils of the locations where leaf litter was collected (Table 1) are characterized by a high availability of Ca (Rubio and Escudero 2005),

which apparently does not limit the tissue lignification of invasive alien species. Although plant species trade off plant growth for tissue lignification (Villar et al. 2006), the elevated C assimilation capacity for growth commonly described in invasive species (Pyšek and Richardson 2007) and the higher levels of litter lignin found in our study, may reflect a C availability sufficient to maximize both strategies. This overall C strategy apparently improves the growth rate and protection of leaf tissues which leads to a higher nutrient use efficiency, another strategy also reported in invasive alien species (Pyšek and Richardson 2007).

Our results do not match those from previous studies comparing leaf litter properties of invasive and native species, as these studies showed that, in general, invasive alien species tended to exhibit higher SLA and higher nutrient concentrations [see Ehrenfeld (2003) for a detailed description]. However, these differences seemed to be biased by the selection of species and regions (mostly N<sub>2</sub>-fixing invasive species in oligotrophic soils; Levine et al. 2003; Yelenik et al. 2007).

### Predictors of litter decomposition rate

N content in litter was related to litter decomposition in all species groups and in both harvests (Table 3). This agrees with the well-known fact that high N content in litter and in

soil promotes decomposition (Ehrenfeld 2003; Cornwell et al. 2008). However, its relative importance was higher in invasive alien species in the final harvest (Table 3), which may be attributed to their higher lignin content. Lignin, which forms complex associations with litter proteins, inhibits the action of decomposing organisms and delays the decomposition processes during the first months (Gallardo and Merino 1993). Thus, N, unlike lignin, becomes an important predictor later on, when soil organism activity overcomes the barrier of the lignin complex and starts to release N from litter proteins (Gallardo and Merino 1992). In addition, P content in leaf litter exhibited a great positive influence on the decomposition of invasive alien litter, but not on native litter (Table 3). Both results are in agreement with Bubb et al. (1998) and Xuluc-Tolosa et al. (2003), who demonstrated that litter with a high lignin content needs a high N content in leaves and soil for its decomposition, and that this process becomes P dependent. Consequently it may become important as a limiting factor for the decomposition of the lignin-rich litter of invasive alien species when lignin degradation by micro-organisms takes off. Perhaps the lower lignin content of the litter of the native species makes them less dependent on P and on N (Table 3; Knorr et al. 2005).

Overall, leaf traits predicted litter decomposition rate better for native than for exotic species, especially at the first harvest. This suggests that other unquantified litter traits of exotic species might initially exert a tighter control on the decomposition process than the measured traits. For instance, high levels of relatively mobile polyphenolic compounds have been commonly reported in invasive species (Ehrenfeld 2003; Knight et al. 2007). These C-based defences may be useful under the full-sun habitat conditions of the early community succession stages, where aliens mainly invade because they provide protection from UV light, and defence against generalist herbivores, but not against the specialist ones (see references in Marko et al. 2008). Such an ecological strategy of invasive species remains in the senesced leaves, influencing the initial processes of litter decomposition (Hättenschwiler and Vitousek 2000).

#### Consequences of invasions for nutrient cycling

Lignin is an immobile C-based substance that influences nutrient cycles at the ecosystem scale by slowing down litter decomposition (Swift et al. 1979; Cornwell et al. 2008). Contrary to most other studies on litter decomposition rates by aliens versus natives, our results have demonstrated that invasive alien species produce less decomposable leaf litter compared to co-familial native Mediterranean plants, probably due to their higher lignin content. Our results, therefore, suggest that the soil nutrient turnover, and thereby

nutrient availability, will not be enhanced upon invasion in Spanish Mediterranean ecosystems; actually they might even be slightly inhibited. However, our experiment has the limitation of not representing natural soil and litter layer conditions in the habitats where invasive species are becoming abundant. In addition, our decomposition bed litter matrix was composed of species (*Quercus* spp.) that generate low decomposition rates (Gallardo and Merino 1993) and possibly select for organisms adapted to the decomposition of lignin-rich litter but reduce soil fauna and microbial diversity (Gallardo and Merino 1992). Thus, changes in decomposer community, soil conditions, or both caused by invasion processes may feedback or moderate the decomposition patterns reported here. For instance, although litters from invasive *Acacia* species are hard to decompose (see “Electronic supplementary material”, Table S1; Yelenik et al. 2007) the N<sub>2</sub>-fixing strategy of these species highly augments N soil availability and microbial soil activity, accelerating litter decomposition upon invasion (Yelenik et al. 2007). High soil N and P availability by natural or human-mediated actions may lead to the same results (Knorr et al. 2005). However, Spanish Mediterranean soils are in general characterized by their low N and P content (Gallardo and Parama 2007) and their low microbial activity and biomass (Gallardo and Merino 1992). Accordingly, a progressive increment of invasive plant species characterized by their lignin-rich litter and without a significant increment in available N and P in soil, would cause a slowing down of nutrient cycling by reducing the litter quality and the abundance and diversity of decomposer communities (Gallardo and Merino 1992; Chigineva et al. 2009). In this sense, Cotrufo et al. (1998) demonstrated that litter of impoverished quality due to an increment in lignin content and lignin:N ratio was avoided by soil fauna in a food choice experiment.

Our results also suggest that under the scenario of a progressive displacement of native species by invasive ones, altering the spectra of growth forms in an ecosystem, invokes the highest relative impact on the litter decomposition rate (the relative impact depending on whether the native community is completely displaced or invasive and native species coexist; Fig. 1). Consequently, changes in community structure from herbaceous to less decomposable invasive woody species, which have been reported for different Mediterranean-type ecosystems worldwide (Di Tomaso 1998; Bousquet-Melou et al. 2005) could augment this negative feedback on litter decomposition. Our data also highlight the importance of Family and, to a lesser extent, Leaf type on the litter decomposition rate (Fig. 1). The impoverishment of taxonomic diversity in native woody communities due to the introduction in forestry of exotic evergreen trees, which belong to a few selected taxonomic groups (e.g. *Pinus*, Pinaceae; *Eucalyptus*, Myrtaceae;

see “Electronic supplementary material”, Table S1 for mean mass loss values), could also augment this negative feedback. There may be worse consequences for nutrient cycling of the invasive plant species on establishment when both effects occur at the same time, as has been reported in the Cape region of South Africa (Manders and Richardson 1992).

In summary, invasive species have the potential to slow down nutrient cycling in Spanish Mediterranean ecosystems through slower litter decomposition. The high lignin content that drives this change is a reflection of the features displayed by alien species which allow them to become invasive. The degree of the impact will be highest when invasive species have a higher lignin content than natives, which include a few genera of woody evergreen alien species.

**Acknowledgements** Thanks are due to Margarita Costa-Tenorio, Antonio Gallardo, Rob Broekman and Manuel Godoy for field help, supplying ion exchange resin membranes and lab assistance, respectively. Financial support was provided by the Spanish Ministry for Education and Science grants RASINV GL2004-04884-C02 02/BOS (as part of the coordinate project RINVE) and CGL2007-61873/BOS.

## References

- Abrams MM, Jarrell WM (1992) Bioavailability index for phosphorous using ion-exchange resin impregnated membranes. *Soil Sci Soc Am J* 56:1532–1537
- Allison SD, Vitousek PM (2004) Rapid nutrient cycling in leaf litter from invasive plants in Hawaii. *Oecologia* 141:612–619
- Armstrong DP, Westoby M (1993) Seedlings from large seeds tolerate defoliation better—a test using phylogenetically independent contrast. *Ecology* 74:1092–1100
- Ashton IW, Hyatt LA, Howe KM, Gurevitch J, Lerdau MT (2005) Invasive species accelerate decomposition and litter nitrogen loss in a mixed deciduous forest. *Ecol Appl* 15:1263–1272
- Bousquet-Melou A, Louis S, Robles C, Greff S, Dupouyet S, Fernandez C (2005) Allelopathic potential of *Medicago arborea*, a Mediterranean invasive shrub. *Chemoecology* 15:193–198
- Bremer B, et al (2003) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG II. *Bot J Linn Soc* 141:399–436
- Bubb KA, Xu ZH, Simpson JA, Saffigna PG (1998) Some nutrient dynamics associated with litterfall and litter decomposition in hoop pine plantations of southeast Queensland, Australia. *For Ecol Manage* 110:343–352
- Burnham KP, Anderson DR (2002) Model selection and multimodel inference. Springer, New York
- Castro-Díez P, González-Muñoz N, Alonso A, Gallardo A, Poorter L (2009) Effects of exotic invasive trees on nitrogen cycling: a case study in Central Spain. *Biol Invasions* 11:1973–1986
- Castroviejo S (1986–2008) Flora Ibérica. Plantas vasculares de la Península Ibérica e Islas Baleares. (todos los vols). Real Jardín Botánico-CSIC, Madrid
- Chigineva NI, Aleksandrova AV, Tiunov AV (2009) The addition of labile carbon alters litter fungal communities and decreases litter decomposition rates. *Appl Soil Ecol* 42:264–270
- Cornelissen JHC (1996) An experimental comparison of leaf decomposition rates in a wide range of temperate plant species and types. *J Ecol* 84:573–582
- Cornwell WK, et al (2008) Plant species traits are the predominant control on litter decomposition rates within biomes worldwide. *Ecol Lett* 11:1065–1071
- Cotruflo MF, Briones MJL, Ineson P (1998) Elevated CO<sub>2</sub> affects field decomposition rate and palatability of tree leaf litter: importance of changes in substrate quality. *Soil Biol Biochem* 30:1565–1571
- Di Tomaso JM (1998) Impact, biology and ecology of saltcedar (*Tamarix* spp.) in the southwestern United States. *Weed Technol* 12:326–336
- Drenovsky RE, Batten KM (2007) Invasion by *Aegilops triuncialis* (barb goatgrass) slows carbon and nutrient cycling in a serpentine grassland. *Biol Invasions* 9:107–116
- Dukes JS, Mooney HA (2004) Disruption of ecosystem processes in western North America by invasive species. *Rev Chil Hist Nat* 77:411–437
- Ehrenfeld JG (2003) Effects of exotic plant invasions on soil nutrient cycling processes. *Ecosystems* 6:503–523
- Epstein E (1994) The anomaly of silicon in plant biology. *Proc Natl Acad Sci USA* 91:11–17
- Fowler J, Cohen L (1992) Practical statistics for field biology. Wiley, London
- Gallardo A, Merino J (1992) Nitrogen immobilization in leaf litter at two Mediterranean ecosystems of SW Spain. *Biogeochemistry* 15:213–228
- Gallardo A, Merino J (1993) Leaf decomposition in two Mediterranean ecosystems of southwest Spain, influence of substrate quality. *Ecology* 74:152–161
- Gallardo A, Parama R (2007) Spatial variability of soil elements in two plant communities of NW Spain. *Geoderma* 139:199–208
- Garnier E, et al (2004) Plant functional markers capture ecosystem properties during secondary succession. *Ecology* 85:2630–2637
- Gasso N, et al (2009) Exploring species attributes and site characteristics to assess plant invasions in Spain. *Divers Distrib* 15:50–58
- Hättenschwiler S, Vitousek PM (2000) The role of polyphenols in terrestrial ecosystem nutrient cycling. *Trends Ecol Evol* 15:238–243
- Hobbie SE (1992) Effects of plant-species on nutrient cycling. *Trends Ecol Evol* 7:336–339
- Knight KS, Kurylo JS, Endress AG, Stewart JR, Reich PB (2007) Ecology and ecosystem impacts of common buckthorn (*Rhamnus cathartica*): a review. *Biol Invasions* 9:925–937
- Knorr M, Frey SD, Curtis PS (2005) Nitrogen additions and litter decomposition: a meta-analysis. *Ecology* 86:3252–3257
- Kolar CS, Lodge DM (2001) Progress in invasion biology: predicting invaders. *Trends Ecol Evol* 16:199–204
- Kueffer C, Klingler G, Zirfass K, Schumacher E, Edwards PJ, Gusewell S (2008) Invasive trees show only weak potential to impact nutrient dynamics in phosphorus-poor tropical forests in the Seychelles. *Funct Ecol* 22:359–366
- Lautner S, Ehrling B, Windeisen E, Rennenberg H, Matyssek R, Fromm J (2007) Calcium nutrition has a significant influence on wood formation in poplar. *New Phytol* 173:743–752
- Levine JM, Vila M, D’Antonio CM, Dukes JS, Grigulis K, Lavorel S (2003) Mechanisms underlying the impacts of exotic plant invasions. *Proc R Soc Lond B Biol Sci* 270:775–781
- Liao CZ, et al (2008) Altered ecosystem carbon and nitrogen cycles by plant invasion: a meta-analysis. *New Phytol* 177:706–714
- Mack RN, Simberloff D, Lonsdale WM, Evans H, Clout M, Bazzaz FA (2000) Biotic invasions: causes, epidemiology, global consequences, and control. *Ecol Appl* 10:689–710
- Manders PT, Richardson DM (1992) Colonization of Cape fynbos communities by forest species. *For Ecol Manage* 48:277–293
- Marko PD, Gross EM, Newman RM, Gleason FK (2008) Chemical profile of the North American native *Myriophyllum sibiricum* compared to the invasive *M. spicatum*. *Aquat Bot* 88:57–65
- Poorter H, Villar R (1997) The fate of acquired carbon in plants: chemical composition and construction costs. In: Bazzaz FA, Grace J

- (eds) Plant resource allocation. Academic Press, San Diego, pp 39–72
- Pyšek P, Richardson DM (2007) Traits associated with invasiveness in alien plants: where do we stand? In: Caldwell MM (ed) Biological invasions, vol 193. Springer, Berlin Heidelberg, pp 97–125
- Pyšek P, Richardson DM, Rejmánek M, Webster GL, Williamson M, Kirschner J (2004) Alien plants in checklist and floras: towards better communication between taxonomist and ecologist. *Taxon* 53:131–143
- Quested HM, et al (2003) Decomposition of sub-arctic plants with differing nitrogen economies: a functional role for hemiparasites. *Ecology* 84:3209–3221
- Quested HM, Callaghan TV, Cornelissen JHC, Press MC (2005) The impact of hemiparasitic plant litter on decomposition: direct, seasonal and litter mixing effects. *J Ecol* 93:87–98
- R Development Core Team (2009) R: a language and environment for statistical computing. R foundation for Statistical Computing, Vienna. ISBN 3-900051-07-0. <http://www.R-project.org>
- Richardson DM, Pyšek P, Rejmanek M, Barbour MG, Panetta FD, West CJ (2000) Naturalization and invasion of alien plants: concepts and definitions. *Divers Distrib* 6:93–107
- Rubio A, Escudero A (2005) Effect of climate and physiography on occurrence and intensity of decarbonation in Mediterranean forest soils of Spain. *Geoderma* 125:309–319
- Sanz Elorza M, Dana Sanchez D, Sobrino Vesperinas E (2004) Atlas de las Plantas Alóctonas Invasoras en España. Ministerio de Medio Ambiente, Madrid
- Sayer EJ (2006) Using experimental manipulation to assess the roles of leaf litter in the functioning of forest ecosystems. *Biol Rev* 81:1–31
- Sims GK, Ellsworth TR, Mulvaney RL (1995) Microscale determination of inorganic nitrogen in water and soil extracts. *Commun Soil Sci Plant Anal* 26:303–316
- Sneller FEC, et al (1999) Toxicity of arsenate in *Silene vulgaris*, accumulation and degradation of arsenate-induced phytochelatins. *New Phytol* 144:223–232
- Soltis DE, et al (2000) Angiosperm phylogeny inferred from 18S rDNA, rbcL, and atpB sequences. *Bot J Linn Soc* 133:381–461
- Strauss SY, Webb CO, Salamin N (2006) Exotic taxa less related to native species are more invasive. *Proc Natl Acad Sci USA* 103:5841–5845
- Subler S, Blair JM, Edwards CA (1995) Using anion-exchange membranes to measure soil nitrate availability and net nitrification. *Soil Biol Biochem* 27:911–917
- Swift MJ, Heal OW, Anderson JM (1979) Decomposition in terrestrial ecosystems. In: *Studies in ecology*, vol 5. Blackwell, Oxford
- Valéry L, Fritz H, Lefeuvre JC, Simberloff D (2008) In search of a real definition of the biological invasion phenomenon itself. *Biol Invasions* 10:1345–1351
- Villar R, Robledo JR, De Jong Y, Poorter H (2006) Differences in construction costs and chemical composition between deciduous and evergreen woody species are small as compared to differences among families. *Plant Cell Environ* 29:1629–1643
- Vitousek PM, Howarth RW (1991) Nitrogen limitation on land and in the sea—how can it occur? *Biogeochemistry* 13:87–115
- Xuluc-Tolosa FJ, Vester HFM, Ramirez-Marcial N, Castellanos-Alboreo J, Lawrence D (2003) Leaf litter decomposition of tree species in three successional phases of tropical dry secondary forest in Campeche, Mexico. *For Ecol Manage* 174:401–412
- Yelenik SG, Stock WD, Richardson DM (2007) Functional group identity does not predict invader impacts: differential effects of nitrogen-fixing exotic plants on ecosystem function. *Biol Invasions* 9:117–125