

Costs *versus* risks: Architectural changes with changing light quantity and quality in saplings of temperate rainforest trees of different shade tolerance

FERNANDO VALLADARES,^{1,2,3*} ALFREDO SALDAÑA⁵ AND ERNESTO GIANOLI^{5,4,6}

¹*Instituto de Recursos Naturales. CCMA. CSIC. Serrano 115 dpdo. E-28006 Madrid. Spain (Email: valladares@ccma.csic.es)*, ²*Departamento de Biología y Geología, Escuela Superior de Ciencias Experimentales y Tecnológicas, Universidad Rey Juan Carlos, c/ Tulipán s/n, Móstoles, Spain;*

³*Laboratorio Internacional de Cambio Global (LINCGlobal), UC-CSIC, Departamento de Ecología, Facultad de Ciencias Biológicas,* ⁴*Center for Advanced Studies in Ecology & Biodiversity (CASEB), P. Universidad Católica, Alameda, Santiago,* ⁵*Departamento de Botánica, Universidad de Concepción, Casilla, Concepción, and* ⁶*Departamento de Biología, Universidad de La Serena, Casilla, La Serena, Chile*

Abstract Light requirements and functional strategies of plants to cope with light heterogeneity in the field have a strong influence on community structure and dynamics. Shade intolerant plants often show a shade avoidance strategy involving a phytochrome-mediated stem elongation in response to changes in red : far red ratio, while shade-tolerant plants typically harvest light very efficiently. We measured plant size, stem diameter, internode and leaf lengths in randomly chosen saplings of 11 woody species differing in their shade tolerance in both a secondary forest and an old-growth temperate evergreen rainforest in southern Chile. We also recorded the irradiance spectrum and the diffuse and direct light availabilities at each sampling point. Significant differences were found for the mean light environment of the saplings of each species, which also differed in basal stem diameter, internode length and leaf length, but not in plant height. Both plant slenderness (plant height/stem diameter) and mean internode length increased with increasing light availability, but no relationship was found between any of these two traits and red : far red ratio. The change in plant slenderness with light availability was of lesser magnitude with increasing shade tolerance of the species, while internode change with light availability increased with increasing shade tolerance of the species. Shade tolerators afford higher costs (thicker stems and plants), which render more biomechanically robust plants, and respond more to the light environment in a trait strongly influencing light interception (internode length) than shade intolerant species. By contrast, less shade-tolerant plants afforded higher risks with a plastic response to escape from the understorey by making thinner plants that were biomechanically weaker and poorer light interceptors. Thus, species differing in their shade tolerances do differ in their plastic responses to light. Our results contribute to explain plant coexistence in heterogeneous light environments by improving our mechanistic understanding of species responses to light.

Key words: elongation, light capture, phenotypic plasticity, red : far red ratio, shade avoidance, shade tolerance, temperate rainforest, understorey.

INTRODUCTION

Plant survival in low light conditions heavily relies on maximization of light interception and utilization for carbon fixation coupled with a minimization of carbon loss by respiration (Valladares & Niinemets 2008). Shade-tolerant plants are also known for their capacity to persist for long periods of time in the understorey, coping with other factors in addition to limiting light such as herbivory and risk of mechanical damage

(Kitajima 1994). While many studies have addressed leaf features conferring shade tolerance (e.g. Craine & Reich 2005; Lusk & Warton 2007), only a few have explored whole-plant features associated with survival in dim understorey light (Reich *et al.* 2003; Baltzer & Thomas 2007). This is remarkable because light interception is a whole crown property (Pearcy *et al.* 2005; Baltzer & Thomas 2007). Light interception by a plant is influenced by a number of traits such as leaf inclination angles, branching patterns and internode lengths, all affecting crown architecture and leaf display (Poorter & Werger 1999; Poorter *et al.* 2003; Valladares & Niinemets 2007). Understorey plants of

*Corresponding author.

Accepted for publication February 2011.

contrasting growth form and shape are known to converge in their light interception efficiency (Valladares *et al.* 2002b), but they do segregate regarding their plastic phenotypic responses to available light (Valladares *et al.* 2000). Shade intolerant plants often exhibit the shade avoidance syndrome (Smith & Whitelam 1997) which involves high morphological and architectural plasticity in order to maximize elongation and apical dominance (Ballare 1999; Weinig 2000). In general, shade-tolerant plants are less plastic than shade intolerant plants (Valladares *et al.* 2002a; Portsmouth & Niinemets 2007), but this typically applies to leaf level traits, and even at the leaf scale, evidence for this is consistent only for certain physiological traits like photosynthetic capacity and stomatal conductance (Valladares & Niinemets 2008). We hypothesized that phenotypic variation of saplings along the light gradient in a forest understorey would reflect interspecific differences in shade tolerance. In particular, shade-tolerant plants are expected to cope with light shortage by affording the costs of producing robust individuals of a high light harvesting efficiency, a conservative strategy that should yield high fitness compared with a shade avoidance syndrome. In contrast, less shade-tolerant plants are expected to show phenotypic variation functional for shade avoidance, investing in vertical growth at the expense of poor light interception and carbon gain, thus risking their survival in the understorey if shading prevails.

There is discrepancy in the relative importance of the specific cue eliciting plastic responses to light (Dorn *et al.* 2000). One group of studies, primarily physiological, emphasizes the phytochrome-mediated responses elicited by changes in the red : far red ratio (R : FR) that is typically low in the dim understorey light (Smith 1995). Another group of studies, primarily ecological, focuses on the responses elicited by the actual intensity of the irradiance and not by its spectral characteristics (e.g. Sánchez-Gómez *et al.* 2006). Recent studies are showing more complex combinations of eliciting cues, so the shade avoidance syndrome might result not only from changes in the light but also in the chemical environment of the plant (Pierik *et al.* 2004; Mullen *et al.* 2006). It has been argued that the phytochrome-mediated responses allow the plant to anticipate competition providing an early neighbour detection (Ballare 1999). But it is the amount of light that eventually determines survival and performance of understorey plants (Dorn *et al.* 2000; Niinemets & Valladares 2006; Sánchez-Gómez *et al.* 2006). Consequently, we hypothesize that traits related with elongation, and thus with shade avoidance, would respond more to R : FR than to light quantity, while traits involved in light harvesting efficiency would respond more to light quantity than to R : FR. These patterns should be evident in the phenotypic variation of saplings along the light gradients

occurring in forest understoreys where light is a key environmental factor driving plant individual performance and community dynamics.

We tested these two groups of hypotheses with a field study of saplings of 11 woody species differing in their shade tolerance in the understorey of the temperate evergreen rainforest in southern Chile. The ecology of these temperate hyper-humid forests is only beginning to be understood but they are ideal for the present study because many plant species coexist in these forests and light availability has been shown to play an important role in vegetation dynamics and niche differentiation (Lusk 1996; Saldaña & Lusk 2003; Lusk *et al.* 2008).

MATERIALS AND METHODS

Study site and sampling

The study took place in the lowland rainforest of Parque Nacional Puyehue (350–440 m a.s.l.; 40°39'S, 72°11'W), located in the western foothills of the Andean range in south-central Chile. Both old-growth and secondary rainforest stands at this altitude are composed exclusively of broad-leaved evergreen tree species (Saldaña & Lusk 2003; Lusk & Piper 2007). The climate is maritime temperate, with an average annual precipitation of around 3500 mm and 13.8°C and 5.4°C mean maximum and minimum temperature, respectively. The dominant canopy species in the mature lowland forest are *Laureliopsis philippiana* (Atherospermataceae), *Aextoxicon punctatum* (Aextoxicaceae), *Nothofagus dombeyi* (Fagaceae) and *Eucryphia cordifolia* (Cunoniaceae). Less abundant canopy trees are *Dasyphyllum diacanthoides* (Asteraceae), *Luma apiculata* (Myrtaceae) and *Weinmannia trichosperma* (Cunoniaceae). Advanced regeneration in the understorey is composed of these tree species in addition to *Amomyrtus luma* (Myrtaceae), *Azara serrata* (Flacourtiaceae), *Caldehuvia paniculata* (Cunoniaceae), *Gevuina avelana* (Proteaceae), *Myrceugenia planipes* (Myrtaceae) and *Rhaphitamnus spinosus* (Verbenaceae) (Saldaña & Lusk 2003). Common woody species in treefall gaps are *Lu. apiculata*, *C. paniculata*, *Embothrium coccineum* (Proteaceae) and *Aristotelia chilensis* (Elaeocarpaceae) (Lusk 2002). In general, most of these tree species have intermediate light requirements (Kneeshaw *et al.* 2006).

Sampling was carried out across mature forest and secondary forest understoreys, which significantly differed in light availability and in the spectral features of the irradiance (Table 1). A minimum of five saplings of the 11 most common woody species were randomly chosen (Table 2).

We recorded for each plant the following morphological traits: sapling height (length of the main stem), basal stem diameter (using a digital calliper with a resolution of 0.01 mm), internode length (average of two consecutive internodes at the middle of the shoot) and the length of leaves associated with the measured internodes. We calculated a slenderness index for each sapling as the plant height : stem diameter ratio. Shade tolerance of each species was assigned from the values of the minimum light requirement given by Lusk *et al.* (2008).

Table 1. Mean values (\pm SE) of light quantity and light quality of gaps, secondary forest and old-growth forest sites in a Chilean rainforest

	Gaps (<i>n</i> = 5)	Secondary forest (<i>n</i> = 14)	Old-growth forest (<i>n</i> = 16)
Light quantity			
Indirect site factor	0.97 \pm 0.01 a	0.20 \pm 0.01 b	0.09 \pm 0.01 c
Direct site factor	0.98 \pm 0.01 a	0.15 \pm 0.01 b	0.06 \pm 0.01 c
Light quality			
Red : far red ratio	1.13 \pm 0.13 a	0.47 \pm 0.02 b	0.26 \pm 0.03 c
Blue : red light ratio	5.84 \pm 0.78 a	6.97 \pm 0.50 b	10.17 \pm 0.47 c
UV-A : red light ratio	0.50 \pm 0.23 a	0.88 \pm 0.15 b	1.92 \pm 0.14 c
UV-B : red light ratio	0.05 \pm 0.04 a	0.06 \pm 0.02 b	0.21 \pm 0.02 c

Values with different letters have significant differences among sites (one-way ANOVA, $P < 0.01$). *n* = sample size.

Table 2. Mean values (\pm SE) of indirect site factor (ISF), direct site factor (DSF), red : far red ratio (R : FR), plant height, basal stem diameter (Basal d.), internode and leaf length of saplings of 11 evergreen species in a Chilean rainforest

Species	<i>n</i>	ISF***	DSF***	R : FR*	Height (cm)	Basal d. (mm)***	Internode (cm)*	Leaf length (cm)***
<i>Aristotelia chilensis</i> (Ac)	6	0.20 \pm 0.02	0.16 \pm 0.02	0.47 \pm 0.07	20.70 \pm 6.79	1.88 \pm 0.66	1.31 \pm 0.51	5.32 \pm 1.91
<i>Amomyrtus luma</i> (Al)	5	0.09 \pm 0.02	0.06 \pm 0.02	0.28 \pm 0.08	25.80 \pm 7.44	3.38 \pm 0.73	1.68 \pm 0.55	3.28 \pm 2.09
<i>Aextoxicon punctatum</i> (Ap)	17	0.16 \pm 0.01	0.12 \pm 0.01	0.34 \pm 0.04	23.50 \pm 4.04	3.29 \pm 0.39	1.82 \pm 0.30	7.81 \pm 1.14
<i>Embothrium coccineum</i> (Ec)	5	0.09 \pm 0.03	0.06 \pm 0.02	0.13 \pm 0.10	45.90 \pm 9.61	6.00 \pm 0.94	1.25 \pm 0.72	10.52 \pm 2.71
<i>Dasyphyllum diacanthoides</i> (Dd)	7	0.09 \pm 0.02	0.07 \pm 0.01	0.24 \pm 0.07	22.94 \pm 6.29	3.11 \pm 0.61	0.93 \pm 0.47	3.03 \pm 1.77
<i>Gevuina avellana</i> (Ga)	8	0.16 \pm 0.02	0.12 \pm 0.01	0.53 \pm 0.06	42.81 \pm 5.88	7.54 \pm 0.58	3.46 \pm 0.44	19.41 \pm 1.66
<i>Luma apiculata</i> (La)	8	0.21 \pm 0.02	0.15 \pm 0.01	0.37 \pm 0.06	17.55 \pm 5.88	1.77 \pm 0.58	1.46 \pm 0.44	2.96 \pm 1.66
<i>Lomatia ferruginea</i> (Lf)	5	0.20 \pm 0.03	0.16 \pm 0.01	0.35 \pm 0.10	14.43 \pm 9.61	3.47 \pm 0.94	1.73 \pm 0.72	7.73 \pm 2.71
<i>Rhaphithamnus spinosus</i> (Rs)	5	0.18 \pm 0.02	0.14 \pm 0.02	0.49 \pm 0.08	27.44 \pm 7.44	3.66 \pm 0.73	1.56 \pm 0.55	2.25 \pm 2.09
<i>Laureliopsis philippiana</i> (Lp)	9	0.14 \pm 0.01	0.10 \pm 0.01	0.26 \pm 0.06	30.73 \pm 5.55	4.36 \pm 0.54	2.56 \pm 0.41	8.47 \pm 1.56
<i>Myrceugenia planipes</i> (Mp)	16	0.15 \pm 0.01	0.11 \pm 0.01	0.36 \pm 0.04	26.55 \pm 4.16	3.56 \pm 0.41	2.12 \pm 0.31	6.36 \pm 1.17

Significant differences among species after one-way ANOVA: * $P < 0.05$, ** $P < 0.01$; *** $P < 0.001$. *n* = sample size.

Determination of light quantity and quality

Light availability over each sapling was quantified by hemispherical photography. Comparisons of methods revealed a good accuracy of hemispherical photography for the description of understory light availability (Bellow & Nair 2003). The photographs were taken using a horizontally levelled digital camera (CoolPix 995, Nikon, Tokyo, Japan), mounted on a tripod and aimed at the zenith, using a fish-eye lens of 180° field of view (FCE8, Nikon). Photographs were analysed for canopy openness using Hemiview canopy analysis software version 2.1 (1999, Delta-T Devices Ltd, UK). This software is based on the program CANOPY (Rich 1990). Photographs were taken under homogenous sky conditions to minimize variations due to exposure and contrast. The direct site factor (DSF) and the indirect site factor were computed by Hemiview accounting for the geographic features of the site. These factors are estimates of the fraction of direct, and diffuse or indirect radiation, respectively, expected to reach the spot where the photograph was taken (Anderson 1966). The hemispheric distribution of irradiance used for calculations of diffuse radiation was standard overcast sky conditions. A total of 160 sky sectors were considered resulting from 8 azimuth \times 20 zenith divisions.

Spectral composition above each sapling of each forest stand was measured using a portable spectroradiometer

(SPEC PAR-NIR, Apogee Inc., Logan, UT, USA), equipped with the cosine corrected sensor with 180° field of view. Spectroradiometer SPEC PAR-NIR has a detection range of 190–2200 nm, with a wavelength accuracy of 0.5 nm. Two days representative of clear-sky conditions (18–19 December 2007) were selected for light-quality measures. We measured light quality at solar midday (12.00–15.00 hours), taking four spectra above each plant at each light environment. The average of these measurements was used as an individual plant value. At the same time we took three spectra in the centre of five large gaps (>2000 m²), as a control of the full sunlight spectral composition. The sequential order of sampling in the field was randomized. Ratio of red to far-red light (R : FR) was estimated by dividing the total irradiance of red light (R: 660 \pm 6 nm wavelength) by the total irradiance of far-red light (FR: 730 \pm 6 nm wavelength). Additionally, we calculated the blue light (450 \pm 6 nm) to red light ratio, UV-A (300 \pm 15 nm) to red light ratio and UV-B (330 \pm 15 nm) to red light ratio.

Data analysis

A one-way ANOVA was used to test for differences in light quantity and quality between forest stands (old-growth




Fig. 1. Irradiance spectra characteristic of gaps and of the understorey of secondary and old-growth forest patches. We included the average spectra of five large gaps (>2000 m²), as a reference for full sunlight spectral composition.

forest, secondary forest and gaps). Species characteristics (light environment and morphological traits) were also compared by a one-way ANOVA. Correlation analyses (Pearson product-moment coefficients) were used to test for the relationship between morphological traits of saplings (slenderness, internode : leaf length ratio and internode length) and light environment conditions (DSF and R : FR). In order to maximize the sample size and depict the general response patterns of plant traits to light quantity and quality, we pooled data from both old-growth and secondary forest stands for correlation analyses.

In order to estimate the phenotypic change of saplings with light quantity and quality, we calculated the slope of the regression between morphological traits and light conditions across the forest (light availability DSF and R : FR) for each species. These species and trait specific slopes were correlated with the minimum light requirement of the species obtained from Lusk *et al.* (2008). MRL is the 10th percentile of the distribution of each species in relation to light availability (% canopy openness), which is an approximation of the lowest light levels tolerated by each species (Lusk *et al.* 2008).

RESULTS

Gaps had not only more light but also a different light quality than that of the two forest understoreys studied (Fig. 1, Table 1). Significant differences were also found for the light environment of the understorey of the two forests. The understorey of the secondary forest had more direct and diffuse light and a higher R : FR than that of the old-growth forest while the reverse was true for blue to red and UV to red light ratios (Table 1, Fig. 2). However, even though mean R : FR was higher in the secondary forest understorey, R : FR was scattered over a wide range of values in both forest types, with values as high as 0.45 in the

doi:10.1111/j.1442-9993.2011.02245.x




Fig. 2. Diffuse irradiance (indirect site factor) (a) and red : far red ratio (b) versus direct irradiance (direct site factor) in the secondary (open symbols) and old-growth (closed symbols) forest plots where the saplings were found.

darkest understorey and as low as 0.09 in the brightest one, with a remarkable overlap in sampling points from the two forests for the 0.1–0.4 range (Fig. 2). Sampling plants over the two types of forest allowed us to have saplings of each species growing over a wide range of light environments differing either in light quantity, light quality or both.

Significant differences were found for the mean light environment of the saplings of the different species, which also differed in basal stem diameter, internode length and leaf length (Table 3). Differences in plant height among species were not significant (Table 3). Both plant slenderness (plant height/stem diameter) and mean internode length increased with increasing light availability when all individuals from the species studied were plot together (Fig. 3). These two variables exhibited high variability for a given light quantity level (r^2 was below 0.1 in the two regressions) due, at least in part, to the different crown architectures of the different species considered. A similarly positive trend was found for most species when regressions were calculated for each species independently (see

Table 3. Range of light environments (direct site factor and red : far red ratio) where saplings of the 11 evergreen species were sampled in a Chilean rainforest

Species	Direct site factor range		Red : far red ratio range	
	Lower value	Upper value	Lower value	Upper value
<i>Aristotelia chilensis</i> (Ac)	0.14	0.19	0.12	0.79
<i>Amyrtus luma</i> (Al)	0.03	0.10	0.23	0.36
<i>Aextoxicon punctatum</i> (Ap)	0.04	0.19	0.12	0.66
<i>Embothrium coccineum</i> (Ec)	0.03	0.10	0.13	0.36
<i>Dasyphyllum diacanthoides</i> (Dd)	0.04	0.08	0.10	0.18
<i>Gevuina avellana</i> (Ga)	0.04	0.19	0.17	0.79
<i>Luma apiculata</i> (La)	0.14	0.19	0.12	0.68
<i>Lomatia ferruginea</i> (Lf)	0.14	0.17	0.12	0.58
<i>Rhaphithamnus spinosus</i> (Rs)	0.04	0.16	0.10	0.75
<i>Laureliopsis philippiana</i> (Lp)	0.03	0.19	0.12	0.68
<i>Myrceugenia planipes</i> (Mp)	0.04	0.19	0.32	0.66




Fig. 3. Slenderness (a and b) and internode length (c and d) of the saplings of the 11 species studied *versus* direct site factor and red : far red ratio in the secondary (open symbols) and old-growth (closed symbols) forest plots where the saplings were found.

[4] Appendix S1–S4). No relationship was found between any of these two traits and R : FR (Fig. 3).

Change of plant slenderness with light availability, estimated from the slope of slenderness *versus* the DSF, increased with increasing light requirements of the species (i.e. with decreasing shade tolerance; Fig. 4). By contrast, change of internode length with light availability, estimated in an analogous way, decreased with increasing light requirements of the species. A similar pattern was found for change of slenderness with R : FR, that is, greater differences

along the light gradient with increasing light requirements of the species, although with greater scatter than that observed for variation with DSF (r^2 of 0.24 *vs.* 0.48, respectively; Fig. 5). No relationship was found between changes in internode length with R : FR and light requirements of the species (Fig. 5).

DISCUSSION

Our study revealed that the phenotypic variation of saplings along the light gradient in a forest understorey was influenced by interspecific differences in shade tolerance. We expected that shade-tolerant plants cope with light shortage by affording the resource costs of producing robust individuals of a high light harvesting efficiency, which should pay off in the long run. Less shade-tolerant plants were expected to show phenotypic variation related to escape from the shade, investing in vertical growth at the expense of poor light interception and carbon gain, which compromises survival in the understorey if shading prevails. In our study, plants from shade-tolerant species increased their robustness (decreased slenderness) in deep shade. This pattern is somewhat coincident with the reported variation of leaf area on a mass basis (SLA), where thicker leaves are found in shade-tolerant species (Lusk & Warton 2007), which is often interpreted in terms of decreasing the risk of premature tissue losses via reduced palatability and fragility of leaves (Kitajima 1994; Lusk 2002). This costly strategy should only pay-off if leaf longevity extends over several years; otherwise the extensive mass and energy investment in aerial plant biomass would be maladaptive. Likewise, the observed robustness of saplings in the shade should be associated with enhanced, long-term survival in the understorey for this strategy to be adaptive.

Plants exposed to reductions in R : FR typically follow the shade avoidance syndrome and elongate




Fig. 4. Mean change in slenderness and internode length with increasing direct irradiance (direct site factor (DSF)) (a and b, respectively) *versus* mean light requirement of each of the 11 woody species studied. See Table 2 for abbreviations of the species.




Fig. 5. Mean change in slenderness and internode length with increasing red : far red ratio (R : FR) (a and b, respectively) *versus* mean light requirement of each of the 11 woody species studied. See Table 2 for abbreviations of the species.

their stems and increase their internode lengths to actively compete for light with neighbours (Smith 2000). But different responses, such as those reported here for shade tolerators, have been described in previous studies, particularly when competition for light is not symmetrical, that is, when it is established between plants of different sizes like trees and understory plants. [6] For instance, Reinhart *et al.* (2006) found a reduction in internode length in *Acer platanoides* seedlings in response to a reduced R : FR, which was interpreted as a strategy to ‘sit and wait’ rather than compete for light with canopy trees (Reinhart 2010). Responses to shade seems to involve a coordinated response to low light intensity and to changes in both blue light and R : FR (Lin 2000; Valladares & Niinemets 2008), which leads, at least in certain plant species, to extend survival in the understory and to increase chances to take advantage of eventual canopy openings.

Interestingly, patterns of phenotypic change arose when light quantity was considered in the analysis, while no relationship was found between plant slenderness and light quality. Shade tolerance was more related to changes in these architectural traits with light quantity than with light quality (R : FR). We hypothesized that traits related with elongation, and thus with shade avoidance, would respond more to R : FR than to light quantity, while traits involved in light harvesting efficiency would respond more to light quantity than to R : FR. Both plant slenderness (plant height/stem diameter) and mean internode length increased with increasing light availability when all individuals from the 11 species studied were plotted together, but no relationship was found between any of these two traits and R : FR. The change in plant slenderness with light availability was of lesser magnitude with increasing shade tolerance of the species, while internode change

1 with light availability increased with increasing shade
2 tolerance of the species.

3 The reduction in internodes lengths of plants
4 experiencing deep shade is a case of counter-gradient
5 variation (Conover & Schultz 1995) because typical
6 intraspecific plastic responses to shade (elongated
7 internodes) oppose interspecific patterns of pheno-
8 typic variation with increasing shade found here
9 (shortened internodes). This can be interpreted con-
10 sidering that elongation in small saplings growing
11 under the deep shade of a tall canopy is a futile and
12 even risky resource investment (Valladares &
13 Niinemets 2008). On the contrary, stem elongation in
14 the secondary forest may be a rewarding strategy if it
15 leads to overtopping competing plants (see also dis-
16 cussion in Henry and Aarssen 2001). Alternatively, the
17 pattern of shortened internodes might be related
18 allometrically with the reduced slenderness syndrome
19 formerly described. The same explanation of a
20 counter-gradient variation could apply to the positive
21 slenderness-light trend observed across species.
22 However, the same trends are found when data are
23 explored within species (see Appendix S1–S4), sug-
24 gesting that this might not be the case in our study
25 system. However, further studies controlling pheno-
26 typic expression of individual genotypes are required
27 to discard this explanation more conclusively.

28 Only light quantity showed a significant association
29 with this pattern of phenotypic variation, suggesting
30 that despite the considerable evidence accumulated on
31 the role of phytochrome-mediated signalling on plant
32 responses to shade, light quality could play a subordi-
33 nate role with regards to light quantity. An earlier study
34 with tropical vine juveniles showed that for two out of
35 three species the reduction in light quantity (photo-
36 synthetic photon flux density) elicited greater
37 responses in plant slenderness and internodes elonga-
38 tion than the reduction in light quality (R : FR) (Lee
39 1988).

40 Although the 11 woody species considered in this
41 study significantly differed regarding their shade toler-
42 ance, the sampled individuals (saplings < 50 cm
43 height) might not represent the shade tolerance of
44 adult trees because this feature is known to change
45 during the ontogeny (Kneeshaw *et al.* 2006; Valladares
46 & Niinemets 2008). Lusk *et al.* (2004) suggested that
47 differences in shade tolerance among evergreens
48 co-occurring in temperate rainforests of southern
49 Chile become increasingly important with increasing
50 plant size and/or age. Interestingly, the effect of size on
51 the light requirements of these plants seems to be
52 related to the successional status of these temperate
53 rainforest species. Species regarded as shade tolerant
54 like *M. planipes*, *A. punctatum* and *L. philippiana*, had
55 low light requirements that did not significantly
56 change across height classes (Donoso 1989; Lusk
57 2002, 2004). By contrast, light requirement of the

58 most shade intolerant species increased at increasing
59 ontogenetic stages. Thus, care must be taken when
60 extrapolating shade tolerance and functional features
61 of saplings growing under low, understory light to
62 older plants and adult individuals even of the same
63 species. Another word of caution is needed regarding
64 the estimated phenotypic variance in response to the
65 light environment; we argue that the variance esti-
66 mated here is meaningful on an ecological basis (i.e.
67 representative according to the co-occurring popula-
68 tions of each species) but this variance might not have
69 captured well the real phenotypic variance of the
70 species, particularly of those having low numbers of
71 individuals in the forest understoreys studied.

72 In conclusion, our study suggests that shade tolera-
73 tors afford higher resource costs (thicker stems and
74 plants), which render more biomechanically robust
75 plants, and respond more to the light environment in
76 traits strongly influencing light interception (internode
77 length in the present study) than shade intolerant
78 species. By contrast, less shade-tolerant plants afforded
79 higher risks in their plastic response to escape from the
80 understory by making thinner plants that are biome-
81 mechanically weaker and poorer light interceptors. Species
82 differing in their shade tolerances differed in their
83 plastic responses to light, more clearly to light intensity
84 than to light quality. Differences in plasticity arose not
85 from overall or mean plasticity but from plasticity for
86 specific traits with contrasting functional meanings.
87 Our results thus contribute to explaining plant coexist-
88 ence in heterogeneous light environments by improving
89 our mechanistic understanding of species responses to
90 light. While some species minimize risks and afford
91 costs, others follow the opposite strategy, and both can
92 render successful. The impact in fitness of following one
93 or another is eventually determined by both the spatial
94 and the temporal grain of environmental heterogeneity,
95 which is highly variable depending not only on the type
96 of ecosystem but also on the structure and dynamics of
97 each particular forest patch. This locally idiosyncratic
98 nature of the light environment further promotes coex-
99 istence at higher scales.

100 ACKNOWLEDGEMENTS 101

102 Thanks are due to L. Corcuera for kind loan of equip-
103 ment and to Mónica Cisternas and Diego Gianoli for
104 help with data collection in the field. Financial support
105 was provided by a collaborative grant between CSIC
106 (Spain) and CONICYT (2007-168), FONDECYT
107 grant 1070503 and by the Spanish grants Consolider
108 Montes (CSD2008_00040), VULGLO (CGL2010-
109 22180-C03-03) and REMEDINAL 2 (CMS2009/
110 AMB-1783). The final stage of manuscript writing
111 was promoted by the CONICYT-ACHA MEC-
112 80090005 project. 113

1 **REFERENCES**

- 2 Anderson M. C. (1966) Stand structure and light penetration. II.
3 A theoretical analysis. *J. Appl. Ecol.* **3**, 41–54.
4 Ballare C. L. (1999) Keeping up with the neighbours: phyto-
5 chrome sensing and other signalling mechanisms. *Trends*
6 *Plant Sci.* **4**, 97–102.
7 Baltzer J. L. & Thomas S. C. (2007) Determinants of whole-
8 plant light requirements in Bornean rain forest tree saplings.
9 *J. Ecol.* **95**, 1208–21.
10 Bellow J. G. & Nair P. K. R. (2003) Comparing common
11 methods for assessing understory light availability in
12 shaded-perennial agroforestry systems. *Agric. Forest*
13 *Meteorol.* **114**, 197–211.
14 Conover D. O. & Schultz E. T. (1995) Phenotypic similarity and
15 the evolutionary significance of countergradient variation.
16 *Trends Ecol. Evol.* **10**, 248–52.
17 Craine J. M. & Reich P. B. (2005) Leaf-level light compensation
18 points in shade-tolerant woody seedlings. *New Phytol.* **166**,
19 710–3.
20 Donoso C. (1989) Antecedentes básicos para la silvicultura del
21 tipo forestall siempreverde. *Bosque* **10**, 37–53.
22 Dorn L. A., Pyle E. H. & Schmit J. (2000) Plasticity to light cues
23 and resources in *Arabidopsis thaliana*: testing for adaptive
24 value and costs. *Evolution* **54**, 1982–94.
25 **10** Henry H. A. L. & Aarssen L. W. (2001) Inter- and intraspecific
26 relationships between shade tolerance and shade avoidance
27 in temperate trees. *Oikos* **93**, 477–87.
28 Kitajima K. (1994) Relative importance of photosynthetic traits
29 and allocation patterns as correlates of seedling shade toler-
30 erance of 13 tropical trees. *Oecologia* **98**, 419–28.
31 Kneeshaw D. D., Kobe R. K., Coates K. D. & Messier C. (2006)
32 Sapling size influences shade tolerance ranking among
33 southern boreal tree species. *J. Ecol.* **94**, 471–80.
34 Lee D. W. (1988) Simulating forest shade to study the develop-
35 mental ecology of tropical plants: Juvenile growth in three
36 vines in India. *J. Trop. Ecol.* **4**, 281–92.
37 Lin C. T. (2000) Plant blue-light receptors. *Trends Plant Sci.* **5**,
38 337–42.
39 Lusk C. H. (1996) Gradient analysis and disturbance history of
40 temperate rain forests of the coast range summit plateau,
41 Valdivia, Chile. *Rev. Chil. Hist. Nat.* **69**, 401–11.
42 Lusk C. H. (2002) Leaf area accumulation helps juvenile ever-
43 green trees tolerate shade in a temperate rainforest. *Oecologia*
44 **132**, 188–96.
45 Lusk C. H. (2004) Leaf area and growth of juvenile temperate
46 evergreens in low light: species of contrasting shade toler-
47 ance change rank during ontogeny. *Fun. Ecol.* **18**, 820–8.
48 Lusk C. H. & Piper F. I. (2007) Seedling size influences rela-
49 tionships of shade tolerance with carbohydrate-storage pat-
50 terns in a temperate rainforest. *Fun. Ecol.* **21**, 78–86.
51 Lusk C. H. & Warton D. I. (2007) Global meta-analysis shows
52 that relationships of leaf mass per area with species shade
53 tolerance depend on leaf habit and ontogeny. *New Phytol.*
54 **176**, 764–74.
55 Lusk C. H., Falster D. S., Jara-Vergara C. K., Jimenez-Castillo
56 M. & Saldaña A. (2008) Ontogenetic variation in light
57 requirements of juvenile rainforest evergreens. *Fun. Ecol.* (in
58 press).
59 **11** Mullen J. L., Weinig C. & Hangarter R. P. (2006) Shade avoid-
60 ance and the regulation of leaf inclination in *Arabidopsis*.
61 *Plant Cell Environ.* **29**, 1099–106.
62 Niinemets U. & Valladares F. (2006) Tolerance to shade, drought
63 and waterlogging of temperate, Northern hemisphere trees
64 and shrubs. *Ecol. Monogr.* **76**, 521–47.
65 Pearcy R. W., Muraoka H. & Valladares F. (2005) Crown archi-
66 tecture in sun and shade environments: assessing function
67 and tradeoffs with a 3-D simulation model. *New Phytol.* **166**,
68 791–800.
69 Pierik R., Cuppens M., Voesenek L. & Visser E. (2004) Inter-
70 actions between ethylene and gibberellins in phytochrome-
71 mediated shade avoidance responses in tobacco. *Plant*
72 *Physiol.* **136**, 2928–36.
73 Poorter L. & Werger M. J. A. (1999) Light environment, sapling
74 architecture, and leaf display in six rain forest tree species.
75 *Am. J. Bot.* **86**, 1464–73.
76 Poorter L., Bongers F., Sterck F. J. & Woll H. (2003) Architec-
77 ture of 53 rain forest tree species differing in adult stature
78 and shade tolerance. *Ecology* **84**, 602–8.
79 Portsmouth A. & Niinemets U. (2007) Structural and physiologi-
80 cal plasticity in response to light and nutrients in five tem-
81 perate deciduous woody species of contrasting shade
82 tolerance. *Fun. Ecol.* **21**, 61–77.
83 Reich P. B., Wright I. J., Cavender-Bares J. et al. (2003) The
84 evolution of plant functional variation: traits, spectra, and
85 strategies. *Int. J. Plant Sci.* **164** (Suppl), S143–S64.
86 Reinhart K. O. (2010) The role of facilitative interactions in tree
87 invasions. *New Phytol.* **187**, 559–62.
88 Reinhart K. O., Gurnee J., Tirado R. & Callaway R. M. (2006)
89 Invasion through quantitative effects: intense shade drives
90 native decline and invasive success. *Ecol. Appl.* **16**, 1821–
91 31.
92 Rich P. M. (1990) Characterizing plant canopies with hemi-
93 spherical photographs. *Remote Sens. Rev.* **5**, 13–29.
94 Saldaña A. & Lusk C. (2003) Influence of overstorey species
95 identity on resource availability and variation in composi-
96 tion of advanced regeneration in a temperate rainforest in
97 southern Chile. *Rev. Chil. Hist. Nat.* **76**, 639–50.
98 Sánchez-Gómez D., Valladares F. & Zavala M. A. (2006) Per-
99 formance of seedlings of Mediterranean woody species
100 under experimental gradients of irradiance and water avail-
101 ability: trade-offs and evidence for niche differentiation. *New*
102 *Phytol.* **170**, 795–806.
103 Smith H. (1995) Physiological and ecological function within
104 the phytochrome family. *Annu. Rev. Plant Physiol. Plant Mol.*
105 *Biol.* **46**, 289–315.
106 Smith H. (2000) Phytochromes and light signal perception by
107 plants – an emerging synthesis. *Nature* **407**, 585–91.
108 Smith H. & Whitelam G. C. (1997) The shade avoidance
109 syndrome: multiple responses mediated by multiple
110 phytochromes. *Plant Cell Environ.* **20**, 840–4.
111 Valladares F. & Niinemets Ü. (2007) The architecture of plant
112 crowns: from design rules to light capture and performance.
113 In: *Functional Plant Ecology* (eds F. Pugnaire & F. Valladares)
114 pp. 11–33. CRC Publishers, 11.
115 Valladares F. & Niinemets Ü. (2008) Shade tolerance, a key
116 plant feature of complex nature and consequences. *Annu.*
117 *Rev. Ecol. Evol. Syst.* **39**, 237–57.
118 Valladares F., Wright S. J., Lasso E., Kitajima K. & Pearcy R.
119 W. (2000) Plastic phenotypic response to light of 16 conge-
120 neric shrubs from a Panamanian rainforest. *Ecology* **81**,
121 1925–36.
122 Valladares F., Chico J. M., Aranda I. et al. (2002a) Greater high
123 light seedling tolerance of *Quercus robur* over *Fagus sylvatica*
124 is linked to a greater physiological plasticity. *Trees Struct.*
125 *Funct.* **16**, 395–403.
126 Valladares F., Skillman J. & Pearcy R. W. (2002b) Convergence in
127 light capture efficiencies among tropical forest understory
128 plants with contrasting crown architectures: a case of mor-
129 phological compensation. *Am. J. Bot.* **89**, 1275–84.
130

1 Weinig C. (2000) Plasticity versus canalization: population dif-
2 ferences in the timing of shade-avoidance responses. *Evolu-*
3 *tion* 54, 441–51.

4
5 **SUPPORTING INFORMATION**

6 Additional Supporting Information may be found in
7 the online version of this article:
8

Appendix S1. Slenderness *versus* direct site factor for
each of the 11 species studied. 9

Appendix S2. Internode length *versus* direct site
factor for each of the 11 species studied. 10

Appendix S3. Slenderness *versus* red : far red ratio
for each of the 11 species studied. 11

Appendix S4. Internode length *versus* red : far red
ratio for each of the 11 species studied. 12

Toppan Best-set Premedia Limited	
Journal Code: AEC	Proofreader: Jason
Article No: 2245	Delivery date: 25 February 2011
Page Extent: 9	Copyeditor: Eleime

AUTHOR QUERY FORM

Dear Author,

During the preparation of your manuscript for publication, the questions listed below have arisen. Please attend to these matters and return this form with your proof.

Many thanks for your assistance.

Query References	Query	Remark
1	AUTHOR: Please check and confirm that the affiliation addresses are correct.	
2	AUTHOR: 'red-far red ratio' has been changed to 'red : far red ratio' throughout for consistency, please confirm that this is OK.	
3	AUTHOR: The word 'understories' has been changed to 'understoreys' (plural form of 'understorey') throughout, please confirm that this is correct.	
4	AUTHOR: 'Supplementary material' has been changed to Supporting information citation 'Appendix S1-S4'. Please confirm that this is correct.	
5	AUTHOR: Please spell out SLA.	
6	AUTHOR: Reinhard <i>et al.</i> 2006 has been changed to Reinhart <i>et al.</i> 2006 so that this citation matches the Reference List. Please confirm that this is correct.	
7	AUTHOR: Henry and Aarssen 2003 has been changed to Henry and Aarssen 2001 so that this citation matches the Reference List. Please confirm that this is correct.	
8	AUTHOR: 'Supplementary material in the Electronic Appendix' has been changed to Supporting information citation 'Appendix S1-S4'. Please confirm that this is correct.	
9	AUTHOR: The word 'phytocrome' has been changed to 'phytochrome', please confirm that this is correct.	
10	AUTHOR: Please provide the volume number for Dorn <i>et al.</i> , 2000.	
11	AUTHOR: If this reference has now been published online, please add relevant DOI information. If this reference has now been published in print, please add relevant volume/page information for Lusk <i>et al.</i> , 2008.	
12	AUTHOR: Please provide the page range and city location of publisher for Valladares & Niinemets, 2007.	
13	AUTHOR: Please confirm that figure part labels ('(a)', '(b)', '(c)' etc.) have been inserted correctly for Figures 2-5.	

14	AUTHOR: As per journal style, table legend should be one, short- to medium-length explanatory sentence, so Table 1 legend has been reworded and column one has been restructured. Please check and confirm that the changes are correct and acceptable.	
----	---	--

UNCORRECTED PROOF